

10/567723


IAPOR Rec'd PCT/PTO 08 FEB 2006

MAILLARD REACTION COMPOUNDS OF CYSTEINE AND A SUGAR HAVING MEATLIKE FLAVOUR

This invention relates to a method of providing meat flavouring to foodstuffs and to novel compounds for providing such flavouring.

5

It is known that the formation of meat flavour proceeds via a sequence of steps known as the Maillard reaction. The sequence may be represented as follows:




10 The Maillard reaction is also known as non-enzymatic browning. The Amadori compounds are key intermediates in the formation of flavours.

One amino acid for which no Amadori compound has hitherto been identified is cysteine ($\text{CH}_2(\text{SH}).\text{CH}(\text{NH}_2).\text{COOH}$). Cysteine plays an important role in the development of flavour in meat, and this is believed to result from its ability to release hydrogen sulphide under the conditions of the Maillard reaction (see, for example, Kobayashi and Fujimaki in Agric. Biol.

5 Chem., 29, 698 81965). Numerous references have attributed to sulphur compounds a central role in the formation of meat flavours. However, unlike all other amino acids, cysteine has hitherto been believed to inhibit the Maillard reaction, because it tends to form relatively stable 2-glycosylthiazolidine-4-carboxylic acids, and this reaction competes with, and may even prevent, the formation of Amadori compounds.

10


It has now been found, surprisingly, that, not only is it possible to make Amadori compounds from cysteine, but also that these compounds are particularly effective at conferring meat flavour in foodstuffs. The invention therefore provides a compound of the formula I:



15

in which R is hydrogen when $n=1-4$, or methyl when $n=0-3$.

Although the compound is depicted as being of formula I, this form of the compound is in fact in equilibrium with two other forms, Ia and Ib, as shown below:



All three forms are included within the scope of this invention. Also included are all other isomeric forms, of which a number are possible (for example, when L-cysteine reacts with a D- or L-sugar).

- 5 The compounds may be prepared by known methods. Amino acids or their salts may be reacted with an aldose either in the presence of about 20% of water (see E.F.L.J. Anet, Austr. J. Chem. 10, 193-197 (1957)) or in an organic solvent (see K. Heyns and H. Paulen, Liebigs Ann. Chem., 622, 160 (1959)). In all preparations of Amadori compounds so far known, a reaction medium with low water content is essential to bring about the required condensation
- 10 with aldoses to glycosyl amines. However, the preparation of the Amadori compounds of cysteine can also conveniently be carried out in dilute aqueous solutions or in mixtures of water with water-miscible solvents. It is not necessary to isolate the initially-formed 2-glycosylthiazolidine-4-carboxylic acids. The formation of the Amadori compounds from cysteine plus sugar or from 2-glycosylthiazolidine-4-carboxylic acid is subject to general acid
- 15 catalysis. For this reason, the reactions are preferably carried out in the presence of buffers having a pH of between 1 and 9, preferably between 2 and 6.

Although the Amadori rearrangement proceeds at room temperature, it is preferred to accelerate it by heating. The preferred temperature will depend on the solvents used and on the

20 reactant concentrations, but the temperature generally lies between 60° and 110°C. At 90°-100°C, the preferred reaction time of the aqueous buffered reaction mixture is from 1-12 hours.

The sugars to be used for the present invention are aldoses with 3-6 carbon atoms, including

25 aldopentoses, aldohexoses and deoxyaldohexoses. Specific sugar examples are the aldopentoses ribose, arabinose, xylose and lyxose, the aldohexoses glucose, mannose and galactose and the 6-deoxyaldohexoses rhamnose, fucose and chinovose. Mixtures thereof can also be used. In addition, all stereoisomeric forms of these sugars may be used.

- 30 The cysteine may be used as the compound itself or in the form of a derivative or a salt, which form free cysteine under the reaction conditions. Both L- and D-cysteine and mixtures thereof

may be used. The molar proportions of the cysteine and the aldose may vary between 2:1 and 1:5, preferably between 1.2:1 and 1:1.5.

The buffers used may comprise mixtures of weak acids and bases or compounds that possess both an acidic and a basic function. Suitable acids include carboxylic acids and phosphoric acid. Typical examples of bases include the salts of the previously-mentioned acids, for example, their alkali metal salts or their salts with nitrogen-containing bases such as pyridine, α -picoline and tertiary amines. Primary and secondary amines may also be used, but preferably only if the reaction medium contains a large excess of water, to prevent the competitive reactions of these amines with the aldose to glycosylamines and Amadori rearrangement compounds of these amines. Examples of compounds with both acidic and basic functions are amino acids or their salts, for example glycine, proline and monosodium glutamate, or ribonucleotides or their salts, such as disodium inosine-5'-monophosphate. The buffers are preferably used in concentrations of from 0.1 to 10 times the molar concentration of cysteine.

The Amadori compounds thus prepared may be isolated as crude products and purified by known methods to give the compounds in colourless, crystalline form.

The Amadori compounds of this invention may be used to impart a meat-like taste to foodstuffs, both meat and non-meat, by incorporation of the compounds or compositions containing them into the foodstuff and subsequent heating. The preferred heating time depends on the water activity, the temperature used and the nature of the Amadori compound, but typically it is about the time needed to fry or cook meat.

25

The invention therefore also provides a method of conferring a meaty flavour or aroma on a foodstuff, comprising the addition to the foodstuff before heating of a compound as hereinabove described. The invention additionally provides the use of a compound as hereinabove described to confer on a foodstuff a meaty flavour or aroma.

30

The concentration required may vary over a wide range, depending on the desired taste and aroma effects, but as a general guide, they are employed in concentrations of from 5 to 5000 ppm by weight of foodstuff in consumable form. Good results are obtained by using from 200-2000 ppm.

5

The Amadori compounds of this invention may be used in combination with other known food additives, such as flavour compounds, aroma enhancers, and the like.

The invention is further described with reference to the following non-limiting examples.

10

EXAMPLE 1

Preparation of Amadori compound of L-cysteine and D-xylose

A solution of 60.5 g (0.50 mol) L-cysteine 82.5 g (0.55 mol) D-xylose and 50 g (about 0.60
15 mol) of a mixture of methylated pyridines in 750 ml demineralised water was heated to about 95°C on a water bath in a 1L round-bottomed flask equipped with a reflux condenser. After 0.5 h, 120 g (2.0 mol) acetic acid was added and the solution was heated for a further 6 h at 90°C. The conversion to Amadori compound was then about 25%. The solution was cooled to room temperature and the solution was then continuously extracted with ether to remove the
20 methylated pyridines. The solution was passed through a 40 x 4 cm column of Bio-Rad AG 50W-X4 (H⁺) cation exchange resin. The column was washed with 1 L water and the absorbed substances were displaced from the column with 0.3N ammonia. Fractions containing more than 50% of the compound were combined and evaporated *in vacuo* to small volume. After the addition of just enough ethanol to cause no cloudiness, crystalline compound separated. The
25 pure compound did not melt but decomposed above 170°C.

Found : C 37.87; H 5.96; N 5.51; S 12.65; O 37.80.

Calc. for C₈H₁₅NSO₆ (253.27) : C 37.94; H 5.97; N 5.53; S 12.66; O 37.90.

The product was dried prior to element analysis (*in vacuo* over P₂O₅, 18 h at 69°C).

30 The results of a ¹³C NMR spectrum (in D₂O; δ in ppm, referenced to sodium 3-trimethylsilyl [2,2,3,3-²H₄]-propionate) were as follows:

6

<u>Major isomer</u>	<u>Minor isomer</u>	<u>assignment</u>
26.9(t)	25.6(t)	C6
53.8(t)	50.2(t)	C3
61.5(d)	57.0(d)	C5
5 65.2(t)	65.2(t)	CH ₂ OH
71.9(d)	71.6(d)	} CHOH
75.0(d)	76.6(d)	
81.5(s)	80.9(s)	C2
173.4(s)	173.0(s)	carboxyl carbon

10

EXAMPLE 2

Preparation of Amadori compound of L-cysteine and D-xylose

L-cysteine.HCl monohydrate (17.6 g; 0.1 mol) was dissolved in 50 ml of demineralized water
 15 contained in a 150-ml round-bottomed flask. Disodium hydrogen phosphate (14.2 g; 0.10 mol)
 or 90% lactic acid (10 g; 0.10 mol) was added and the pH of the resulting solution adjusted to
 5.0 by slowly adding with stirring an aqueous 50% sodium hydroxide solution. The sugar
 (0.11 mol) was then added and the mixture stirred for 1 h at 50 °C to obtain an equilibrium
 mixture of thiazolidine, cysteine and sugar. After cooling to room temperature, the pH was re-
 20 adjusted to 5.4 by addition of 50% sodium hydroxide solution. The water content was adjusted
 by evaporation *in vacuo* to 30%. The flask was then equipped, via a double neck adapter, with
 a mechanical stirrer and a reflux condenser, and heated with stirring for 7 h in a thermostated
 water bath of 80 °C. The conversion to Amadori compound is 35%. Isolation was performed
 as described in Example 1.

25

EXAMPLE 3

Continuous process for production of Amadori compound of L-cysteine and D-xylose

The continuous reactor consisted of a 45x2.5 ID cm glass column with temperature control
 30 jacket packed with Bio-Rex 9 anion exchange resin (OH⁻), 100-200 mesh. A restriction
 consisting of a 500 x 0.3 mm ID TeflonTM tubing was connected to the column outlet to
 prevent the formation of gas bubbles in the column. The column was thermostated at 90 °C. A

50% thiazolidine solution, prepared by heating equimolar amounts of cysteine and xylose in water for 1 h at 50 °C, was pumped through the column at a rate of 0.5 ml/min. Analysis of samples taken at regular time intervals showed that the conversion to Amadori compound increased from 12% after 2 hrs to 35% after 6 hrs. The yield of the corresponding batch process was 22.5%. The Amadori compound was isolated from the column eluate by
 5 *evaporation in vacuo* and crystallization of the residue from water-ethanol as in Example 1.

EXAMPLE 4

Preparation of the Amadori compound of L-cysteine and D-ribose

10

The same equipment was used as was used in Example 1. 12.1g (0.1 mol) of L-cysteine, 15g (0.1 mol) of D-ribose and 27.2g (0.2 mol) of potassium dihydrogen phosphate in 150 ml demineralised water were heated for 6h at ca. 90°C under nitrogen. The resulting brown solution was passed through a 35 x 2.5cm column of DowexTM 50W-X4 (H⁺) cation exchange
 15 resin. The neutral components were washed from the column with 1L water and the nitrogen-containing substances were displaced from the column with 0.3N ammonia. Fractions containing more than 50% of the compound were combined and evaporated *in vacuo* to small volume. After the addition of just enough ethanol to cause no cloudiness, crystalline compound separated. The pure compound did not have a sharp melting point but decomposed
 20 at 132°-135°C. The results of a ¹³C NMR spectrum (in D₂O; δ in ppm) were as follows:

	<u>Major isomer</u>	<u>Minor isomer</u>	<u>assignment</u>
	26.8(t)	25.7(t)	C6
	55.2(t)	51.1(t)	C3
25	61.9(d)	57.3(d)	C5
	65.6(t)	65.3(t)	CH ₂ OH
	74.0(d)	73.4(d)	} CHO
	75.9(d)	77.8(d)	
	82.7(s)	81.1(s)	C2
30	174.0(s)	173.5(s)	carboxyl carbon

EXAMPLE 5

Preparation of the Amadori compound of L-cysteine and L-arabinose

The process of Example 3 was repeated with L-arabinose instead of D-xylose. The resulting
5 crystalline material melted with decomposition at 171-173 °C.

The results of a ^{13}C NMR spectrum (in D_2O ; δ in ppm) were as follows:

	<u>Major isomer</u>	<u>Minor isomer</u>	<u>assignment</u>
	26.9(t)	25.8(t)	C6
10	54.9(t)	51.7(t)	C3
	62.1(d)	57.2(d)	C5
	65.4(t)	65.6(t)	} CH ₂ OH CHOH
	73.7(d)	73.9(d)	
	77.7(d)	76.2(d)	
15	81.1(s)	82.8(s)	C2
	174.0(s)	173.6(s)	carboxyl group

EXAMPLE 6

Preparation of the Amadori compound of L-cysteine and D-glucose

20

An aqueous solution of 12.1g (0.1 mol) of L-cysteine, 19.8g (0.1 mol) of D-glucose and 11.5g (0.1 mol) of L-proline was concentrated *in vacuo* to a syrup containing approximately 20% of water. The syrupy mixture was heated on a water bath at 95°C for 2.5h. HPLC analysis showed that, in addition to the Amadori compound of L-cysteine, the Amadori compound of
25 L-proline was also formed, in the ratio of 3:1. The brown reaction mixture was dissolved in 500ml water and the Amadori compounds recovered as described in Example 2. The results of a ^{13}C NMR spectrum (in D_2O ; δ in ppm) were as follows:

	<u>Major isomer</u>	<u>Minor isomer</u>	<u>assignment</u>
	27.1(t)	25.9(t)	C6
	54.1(t)	50.3(t)	C3
	61.6(d)	57.2(d)	C5
5	65.1(t)	65.1(t)	C6'
	63.3(t)	63.0(t)	CH ₂ OH
	71.6(d)	71.3(d)	} CHO
	73.3(d)	73.5(d)	
	74.4(d)	76.1(d)	
10	81.8(s)	81.1(s)	C2
	173.6(s)	173.0(s)	carboxyl carbon

EXAMPLE 7

Preparation of the Amadori compound of L-cysteine and L-rhamnose

15

A solution of 3.3 g (0.30 mol) L-cysteine 49g (0.27 mol) L-rhamnose monohydrate and 30 g (about 0.0 mol) of a mixture of methylated pyridines and 72g (1.20 mol) acetic acid in 450 ml demineralised water was heated for 8h on a steam bath in a 1L round-bottomed flask equipped with a reflux condenser. The resulting brown reaction mixture was purified as described in

20 Example 1. The resulting crystalline compound decomposed at 183°C. The results of a ¹³C NMR spectrum (in D₂O; δ in ppm) were as follows:

10

	<u>Major isomer</u>	<u>Minor isomer</u>	<u>assignment</u>
	21.1 or 21.2(q)	21.1 or 21.2(q)	CH ₃
	26.9(t)	25.9(t)	C6
	54.3(t)	50.8(t)	C3
5	62.1(d)	57.0(d)	C5
	69.9(d)	69.7(d)	
	75.2(d)	74.7(d)	CHOH
	75.8(d)	75.4(d)	
	81.3(s)	81.8(s)	C2
10	173.9(s)	173.5(s)	carboxyl carbon

EXAMPLE 8

Sensory evaluation of Amadori compounds

15 A beef broth was prepared. Four portions of this broth were treated as follows:

- Portion A: no treatment (control);
- Portion B: addition of 480ppm (4.0mmol/L) L-cysteine + 552ppm (4.0 mmol/L) D-xylose;
- Portion C: addition of 1000ppm (4.0 mmol/L) 2-xylosylthiazolidine-4-carboxylic acid
 20 (prepared according to R. Bognár *et al*, Liebigs Ann.Chem.738, 68 (1970);
- Portion D: addition of 1000ppm (4.0 mmol/L) of the Amadori compound of Example 1.

The mixtures were bottled and sterilised in an autoclave for 1h at 100°C. On sensory evaluation, Mixture D was preferred, because of its much stronger meaty taste. Moreover, it
 25 lacked the cysteine-like off-taste of Mixture B and the hydrogen sulphide-like off-odour of Mixture C.

EXAMPLE 9

Test in a gravy

A gravy was prepared by heating 100g of commercial, flavoured margarine to 160°C. On
5 reaching this temperature, 35ml water was added. This was used as an unflavoured control.


A second gravy was prepared by adding 100mg (1000ppm) of the Amadori compound of
Example 2 to 100g of margarine and heating to 160°C. When this temperature was reached,
35ml water was added.

10

A taste comparison revealed a considerable preferment for the gravy containing the Amadori
compound, because of its pronounced, nicely roast, meaty taste.

CLAIMS

1. A compound of the formula I:



5

in which R is hydrogen when $n=1-4$, or methyl when $n=0-3$.

2. A method of preparing a compound according to claim 1, comprising the reaction of cysteine with a sugar in either aqueous solution or in a mixture of water and water-miscible solvents.
3. A method according to claim 2, in which the reaction is accelerated by heating, preferably to between 60° and 110°C .
4. A method according to claim 2, in which the reaction is carried out in the presence of a buffer having a pH of from 1-9, preferably of from 2-6.
5. A method of conferring a meaty flavour or aroma on a foodstuff, comprising the addition to the foodstuff before heating of a compound according to claim 1.
6. Use of a compound according to claim 1 to confer on a foodstuff a meaty flavour or aroma.

20

25

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07C323/58 A23L1/231

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07C A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3 620 772 A (KITADA NAGAYOSHI ET AL) 16 November 1971 (1971-11-16) the whole document -----	1-6
X	GB 1 232 719 A (AJINOMOTO CO.) 19 May 1971 (1971-05-19) the whole document -----	1-6
X	GB 1 469 331 A (PFIZER LTD) 6 April 1977 (1977-04-06) the whole document -----	1-6
X	GB 836 694 A (UNILEVER LTD) 9 June 1960 (1960-06-09) the whole document -----	1-6
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- * & * document member of the same patent family

Date of the actual completion of the international search

6 October 2004

Date of mailing of the international search report

07/12/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Lorenzo Varela, M.J.

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 592 917 A (TANDY JOHN S) 3 June 1986 (1986-06-03) the whole document	1-6
X	US 3 741 775 A (LEE C) 26 June 1973 (1973-06-26) the whole document	1-6
X	US 3 532 514 A (MAY CHARLES GERARD) 6 October 1970 (1970-10-06) the whole document	1-6
X	US 3 660 114 A (THOMAS PAUL D) 2 May 1972 (1972-05-02) the whole document	1-6
X	GB 1 082 504 A (PFIZER & CO C) 6 September 1967 (1967-09-06) the whole document	1-6
X	US 5 723 504 A (MIODUSZEWSKI JAN ZBIGNIEW ET AL) 3 March 1998 (1998-03-03) the whole document	1-6
X	GB 1 285 568 A (UNILEVER LTD.) 16 August 1972 (1972-08-16) the whole document	1-6
X	EP 0 320 057 A (UNILEVER PLC ; UNILEVER NV (NL)) 14 June 1989 (1989-06-14) the whole document	1-6

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3620772	A	16-11-1971	FR 2005135 A5	05-12-1969
			GB 1206265 A	23-09-1970
			IT 1045511 B	10-05-1980
GB 1232719	A	19-05-1971	CH 503470 A	28-02-1971
			FR 2012745 A5	20-03-1970
GB 1469331	A	06-04-1977	BE 825417 A1	11-08-1975
			CA 1047306 A1	30-01-1979
			DE 2506715 A1	21-08-1975
			DK 57175 A	27-10-1975
			FR 2260957 A1	12-09-1975
			IE 40609 B1	04-07-1979
			JP 50111265 A	01-09-1975
			LU 71861 A1	05-01-1977
			NL 7501741 A	20-08-1975
			US 3966985 A	29-06-1976
GB 836694	A	09-06-1960	NONE	
US 4592917	A	03-06-1986	AT 35372 T	15-07-1988
			AU 572626 B2	12-05-1988
			AU 3956685 A	24-10-1985
			CA 1233692 A1	08-03-1988
			DE 3563506 D1	04-08-1988
			EP 0160794 A1	13-11-1985
			ES 8608294 A1	01-12-1986
			GB 2157538 A ,B	30-10-1985
			HK 4989 A	27-01-1989
			JP 60232073 A	18-11-1985
			KR 9205996 B1	25-07-1992
			NZ 211228 A	30-10-1987
			PH 21430 A	15-10-1987
			ZA 8501457 A	30-10-1985
US 3741775	A	26-06-1973	NONE	
US 3532514	A	06-10-1970	GB 1130631 A	16-10-1968
			DE 1692794 A1	03-06-1971
			FR 1470924 A	24-02-1967
			IT 1050403 B	10-03-1981
			NL 6603250 A ,B	12-09-1966
			SE 341316 B	20-12-1971
			SE 362004 B	26-11-1973
			US 3493395 A	03-02-1970
US 3660114	A	02-05-1972	AT 290272 B	25-05-1971
			BE 713352 A	08-10-1968
			CH 497138 A	15-10-1970
			DE 1767138 B1	04-05-1972
			ES 352535 A1	16-07-1969
			FR 1569864 A	06-06-1969
			GB 1182976 A	04-03-1970
			NL 6805127 A	14-10-1968
GB 1082504	A	06-09-1967	AT 273636 B	25-08-1969
			BE 670702 A	08-04-1966
			CH 468157 A	15-02-1969

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 1082504	A		DE 1517100 A1	11-12-1969
			FR 1450072 A	06-05-1966
			IT 1008502 B	30-11-1976
			NL 6513067 A	12-04-1966
US 5723504	A	03-03-1998	PL 293464 A1	23-08-1993
			AT 187734 T	15-01-2000
			AU 3496693 A	03-09-1993
			BG 62172 B1	30-04-1999
			BG 98956 A	28-07-1995
			BR 9305878 A	19-08-1997
			CA 2130106 A1	14-08-1993
			CN 1075969 A	08-09-1993
			CZ 9401945 A3	15-12-1994
			DE 69327310 D1	20-01-2000
			DE 632813 T1	15-02-1996
			WO 9316087 A2	19-08-1993
			EP 0632813 A1	11-01-1995
			ES 2072244 T1	16-07-1995
			FI 943733 A	12-08-1994
			GR 95300017 T1	30-04-1995
			HU 70434 A2	30-10-1995
			JP 7506344 T	13-07-1995
			MX 9300759 A1	01-11-1993
			NO 942930 A	08-08-1994
			PL 171319 B1	30-04-1997
			RO 115633 B1	28-04-2000
			RU 2141337 C1	20-11-1999
			SG 52390 A1	28-09-1998
			SK 86994 A3	08-02-1995
GB 1285568	A	16-08-1972	BE 747962 A1	25-09-1970
			CH 570120 A5	15-12-1975
			DE 2014477 A1	28-01-1971
			ES 377811 A1	16-10-1972
			FR 2039923 A5	15-01-1971
			IT 1050422 B	10-03-1981
			NL 7004242 A	29-09-1970
			ZA 7001973 A	24-11-1971
EP 0320057	A	14-06-1989	AT 88063 T	15-04-1993
			CA 1339870 C	19-05-1998
			DE 3880278 D1	19-05-1993
			DE 3880278 T2	16-09-1993
			EP 0320057 A2	14-06-1989
			ES 2041304 T3	16-11-1993
			JP 1206968 A	21-08-1989
			US 5045335 A	03-09-1991